

The fish fauna of streams in the Madeira-Purus interfluvial region, Brazilian Amazon

Daniela de França Barros^{1*}, Jansen Zuanon², Fernando Pereira de Mendonça³, Helder Mateus Viana Espírito Santo⁴, André Vieira Galuch² and Ana Luisa Mangabeira Albernaz⁵

- 1 Universidade Federal do Pará, Pós-Graduação em Ciências Ambientais. Rua Augusto Corrêa, 1, Guamá. CEP 66075-110. Belém, PA, Brazil.
 - 2 Instituto Nacional de Pesquisas da Amazônia, Coordenação de Pesquisas em Biodiversidade. Avenida André Araújo, 2936, Caixa Postal 478. CEP 69060-001. Manaus, AM, Brazil.
 - 3 Universidade Federal do Amazonas, Instituto de Saúde e Biotecnologia. Coari-Mamiá, 305, Espírito Santo. CEP 69460-000. Coari, AM, Brazil.
 - 4 Instituto Nacional de Pesquisas da Amazônia, Pós-Graduação em Ecologia. Avenida André Araújo, 2936, Aleixo, Caixa Postal 478. CEP 69060-001. Manaus, AM, Brazil.
 - 5 Museu Paraense Emílio Goeldi, Coordenação de Ciências da Terra e Ecologia. Avenida Perimetral, 1901. CEP 66077-830. Belém, PA, Brazil.
- * Corresponding author. E-mail: dani_fbarros@yahoo.com.br

ABSTRACT: Small headwaters streams of the Neotropical region usually have high species richness and diversity. This study aimed to investigate the species composition and abundance of fish fauna in the headwaters streams of the Madeira-Purus interfluvial plain in the Brazilian Amazon. A total of 22 streams of 1st to 3rd order were sampled during two expeditions at two separate locations in April-May and July of 2007. A total of 5508 fishes were captured using hand and small seine nets, belonging to 78 species, 22 families and six orders. Characiformes was the most diverse taxonomic group in the samples, followed by Gymnotiformes and Siluriformes. Our findings indicate that the fish fauna of streams in the Madeira-Purus interfluvial plain is both rich and diverse and should be considered during the implementation of strengthened environmental conservation strategies in this region.

INTRODUCTION

The Neotropical region harbors a high diversity of freshwater fish, which includes almost 4500 already described species and nearly 1550 species still waiting for a scientific description (Reis *et al.* 2003). The Amazon basin is considered to shelter most of this diversity; however knowledge about the ichthyofaunal composition of this region is concentrated in the most accessible areas near large urban centers (Lowe-McConnell 1999). Furthermore, information about fish has been obtained mostly from the main tributaries of the Amazon River, and primarily targeting medium- to large-sized species of commercial value (Böhlke *et al.* 1978; Rapp Py-Daniel and Leão 1991).

Small streams, of the 1st (no tributaries), 2nd (two first order streams join) and 3rd order (two second order streams join), have a unique species pool, and contribute significantly to regional diversity (*e.g.* Espírito-Santo *et al.* 2009). Furthermore, because of the isolated condition of the streams, they are considered important for evolutionary processes, such as speciation (Barthem 2004). However, studies in these environments are limited because of their remoteness (and hence difficult access) and because they shelter small bodied species that are of no commercial interest for use as human food (Lowe-McConnell 1999).

The interfluvial plain between the Madeira and Purus Rivers is one of the most pristine parts of the Amazonian region, but is also one of the least studied with a small amount of protected area coverage. In addition, the Brazilian Federal Government development policies for this region include paving an existing road and the opening of hydroways, dam construction and gas transportation

systems. The implementation of such actions is likely to have a major impact to the region, through the encroachment of human settlements, pollution, and deforestation (Fearnside and Graça 2006). Hence there is serious concern for the conservation of small streams and forested environments, which are particularly vulnerable to human occupation/activities and its impacts (*e.g.* Dias *et al.* 2009).

This study presents the first standardized survey conducted in the interfluvial plain of the Madeira and Purus Rivers. The study aims to contribute to existing knowledge of the fish fauna of small Neotropical streams, comparing our findings to similar studies in other regions in the Brazilian Amazon, and consider the conservation importance of this remote area of the basin.

MATERIALS AND METHODS

Study area

The study was conducted in the area surrounding the BR-319 highway (connecting Porto Velho, in Rondonia state, to Manaus, in the Amazonas state), between the Madeira and Purus Rivers, in southwestern Brazilian Amazon. Two sites were selected for data collection: the first in the southern part of the interfluvial plain near to Porto Velho (Rondônia state) and the second in the northern part, near to Manaus municipality (Amazonas state).

A total of 22 small streams of 1st to 3rd order were sampled, in two separate expeditions; 11 during the first trip between April and May 2007 to the southern part of the region, and 11 streams during the second trip in July 2007 in the northern part of the region. In the first expedition all sampled streams were in close proximity

to site #1 and belong to the Purus river basin, and in the second expedition three streams were sampled near site #2, four near site #3, and four near site #4, all draining into the Madeira river (Figure 1). The duration of each expedition was approximately 20 days. Both expeditions were conducted at the beginning of the dry season with the southern part being visited first due to the rain stopping earlier than at the northern part.

The geology of the region is primarily comprised by Cenozoic deposits, with sediments from the Tertiary and early Quaternary periods, which make up the Solimões Formation. The relief is classified as one of the morpho-structural units of the “lowered Amazonian tableland” (Radambrasil 1978). Soils are mostly red-yellow podzols, but there are also patches of red-yellow latosols, yellow latosols and hydromorphic laterites (Radambrasil 1978).

The climate is equatorial, warm and humid, including the categories of ‘tropical monsoon’ (Am), ‘tropical rainforest’ (Af), and ‘tropical wet and dry’ (Aw) based on the Köppen climate classification system (Radambrasil 1978). The mean annual temperatures ranges between 24 °C and 26°C. The wet season usually starts in October, with the highest volume of rain falling from January to March. Vegetation cover is dominated by dense and open tropical forests (Radambrasil 1978).




FIGURE 1. Study area at the interfluvial region of Madeira and Purus Rivers (dark gray) in the Brazilian Amazon biome (light gray). The numbers indicate the sampling sites at the southern (site #1, 11 streams sampled; Purus basin, near to Porto Velho, Rondônia state) and northern (sites #2, three streams; #3, four streams; and #4, four streams; Madeira basin, near to Manaus, Amazonas state) portions of the interfluvial plain.

Fish sampling

The fish sampling protocol used in this study follows Mendonça *et al.* (2005), for the purposes of comparative analysis with similar studies (*e.g.* Dias *et al.* 2009; Espírito-Santo *et al.* 2009). In each stream that was surveyed, a 50 m long section was delimited for the collection of fish. The two extremities of the section were blocked using a fine-mesh net (5 mm between opposite knots) with the aim to prevent the escape of fishes during capture. Fish were captured using a purse seine of 2 m length (mesh size= 2 mm) and hand nets. Capture was conducted during daylight hours only and always in the same direction; from downstream to upstream. Sampling effort was standardized as much as possible with three collectors working for a 2 h period in each 50 m section.

The collected specimens were sacrificed in a solution of clove oil (Eugenol, 2 drops per litre; *cf.* American Veterinary Medical Association 2001), and fixed in 10% formalin solution. All fish collected in this study were transported to the Instituto Nacional de Pesquisas da Amazônia (INPA) in Manaus. The fish species were separated, and identified by the research team, with the aid of taxonomic literature, and identification keys (*e.g.* Weitzman 1960; 1978; Géry 1977; 1993; Kullander 1986; 1989; 1995; Vari and Ortega 1986; Lucena 1987; Weitzman and Vari 1987; Burgess 1989; Kullander and Ferreira 1991; Huber 1992; Vari 1992; Buckup 1993; Silfvergrip 1996; Reis 1997; Schaefer 1997; Zarske and Géry 1997; Lima and Toledo-Piza 2001; Römer 2002; Crampton *et al.* 2003; Crampton and Albert 2003; Costa 2004; Crampton *et al.* 2004; Crampton and Albert 2004; Mago-Leccia 1994; Crampton *et al.* 2005; Favorito *et al.* 2005; Lundberg 2005; Reis *et al.* 2005; Sousa and Py-Daniel 2005; Thacker *et al.* 2006; Zarske and Géry 2006; Rocha *et al.* 2008; Sarmento-Soares and Martins-Pinheiro 2008; Oyakawa and Mattox 2009; Marinho and Langeani 2010). Samples of the collected specimens were subsequently deposited in the INPA Fish Collection (Table 1). Fishes were collected under the IBAMA license # 10199-1.

RESULTS AND DISCUSSION

A total of 5508 fishes were collected belonging to 78 species, 22 families and six orders. The distribution of most species was related to drainage basins. Of the total species collected, 25 were found in both basins, 19 were collected only in the Purus basin, and 34 only in the Madeira basin (Table 1). Of the 78 species, 12 were identified to the genus level only, with an additional three requiring confirmation of identification to the species level. In other words, the taxonomic status of 19.2% of the collected species was undefined and may possibly be revealed as new species following review by taxonomists.

Characiformes was the most diverse and abundant order, with 39 species and 81.6% of all collected specimens. Based on species number, the second most represented order was Gymnotiformes ($n = 15$), followed by Siluriformes ($n = 12$), Perciformes ($n = 8$), Cyprinodontiformes ($n = 2$), and Synbranchiformes ($n = 2$). Based on sample abundance, the second most represented order was again Gymnotiformes (6.3% of sampled specimens), followed by Perciformes (4.8%), Siluriformes (4.2%), Cyprinodontiformes (2.7%), and Synbranchiformes (0.4%). Furthermore, three families represented almost 80% of sampled individuals (Table 1), comprising Characidae with 56.4% of the total abundance, followed by Lebiasinidae (18.1%), and Crenuchidae (5.2%).

Overall, five species comprised 56.7% of the total abundance; these included *Hyphessobrycon heterorhabdus* Ulrey, 1894 (20.4%), *Hemigrammus vorderwinkleri* Géry, 1963 (11.8%), *Copella nattereri* (Steindachner, 1876) (11.3%), *Hemigrammus schmardae* (Steindachner, 1882) (6.9%), and *Hemigrammus belottii* (Steindachner, 1882) (6.3%). A further six species were found in over 60% of the sampled streams (Table 1); these included *Apistogramma agassizi* (Steindachner, 1875) (77.3%), *Hyphessobrycon heterorhabdus* (Ulrey, 1894) (72.7%) and *Pyrrhulina brevis* Steindachner, 1876 (72.7%), *Rivulus* sp. (68.2%),

Erythrinus erythrinus (Bloch and Schneider, 1801) (63.6%) and *Gymnotus coropinae* Hoedeman, 1962 (63.6%).

Existing research has indicated a large diversity in South American freshwater fish fauna, with Characiformes and Siluriformes usually being most dominant (Castro 1999; Lowe-McConnell 1999; Reis et al. 2003; Buckup et al. 2007). These observations have been further supported by studies in a number of Brazilian Amazon streams (Sabino and Zuanon 1998; Araújo-Lima et al. 1999; Bührnheim and Cox-Fernandes 2001; Dias et al. 2009; Espírito-Santo et al. 2009; Oliveira et al. 2009). In this study, Characiformes was also the most diverse order. However, the second order in species richness was Gymnotiformes, rather than Siluriformes as was found in the the above mentioned studies. In this study, Gymnotiformes represented both a relatively high proportion of species (19.2%), and a relatively high abundance (6.3%). The reason for this may have been due to the high proportion of litter banks in sampled streams, a substrate on which individuals of this order were collected. The species of this order usually inhabit slow moving or lentic (i.e. lake, pond or swamp like) environments (Fink and Fink 1978), which explain our results. On the other hand, the Perciformes, Cyprinodontiformes and Synbranchiformes all represented small proportions of the overall species abundance and richness, which was similar to that found in previous studies (Silva 1995; Mendonça et al. 2005; Espírito-Santo et al. 2009; Oliveira et al. 2009) and other freshwater environments of South America (Lowe-McConnell 1999).

The present study found very high fish species richness

(78 spp.) when compared to existing studies of streams in the Brazilian Amazon (e.g. Araújo-Lima et al. 1999; Bührnheim and Cox-Fernandes 2001). For example, in the studies by Mendonça et al. (2005) and Espírito-Santo et al. (2009), in which 38 sections and 31 sections of creeks of 1st to 3rd order were sampled respectively, both following the same methodology used in this study, a total of 49 and 53 fish species were detected respectively. In addition to the relatively high species richness recorded in this study, there is other evidence of high species diversity in the Madeira-Purus interfluvial region, and hence its conservation importance. Field studies in the Amazonian region have uncovered about 4-5% of previously unnamed species (Peres 2005), which is much smaller than the 19.2% of species with undefined taxonomic status found in this study. Further analysis of the fish collected in this study, that do not match species found in biological collections and/or the specialised literature, may result in the classification of new species. Such species are likely to be of restricted distribution, enlarging the list of endemic fish proposed by Nogueira et al (2010). In conclusion, the results of this study reinforce the requirement for more targeted surveys on the distribution of Amazonian freshwater fish fauna in remote areas. The acquisition of such scientific information is essential to improve knowledge on distribution patterns of fish species; to monitor the impacts of anthropogenic activities, which can possibly threat species survivorship (Peres et al. 2010); and for implementation of effective conservation strategies for ecosystems supporting the fish fauna of Neotropical streams.

TABLE 1. Composition of the fish fauna sampled in 22 streams of the Madeira-Purus interfluvial plain (Rondônia and Amazonas states, Brazilian Amazon). Purus/Madeira= number of individuals captured in each drainage basin; N = total abundance; FR%= relative frequency of the species in relation to the total number of collected specimens; FO%= frequency of occurrence in the sampled streams.

TAXON	BASINS		N	FR (%)	FO (%)	INPA #
	PURUS	MADEIRA				
CHARACIFORMES						
Characidae						
<i>Amazonspinther dalmata</i> Bührnheim, Carvalho, Malabarba and Weitzman, 2008	8	4	12	0.22	18.2	28659, 28660, 28662, 28663, 28665
<i>Aphyocharacidium bolivianum</i> Géry, 1973	0	52	52	0.94	4.5	35819
<i>Bario steindachneri</i> (Eigenmann, 1893)	3	0	3	0.05	9.1	35772, 35784
<i>Bryconops aff. magoi</i> Chernoff and Machado-Allison, 2005	28	0	28	0.51	9.1	35801
<i>Charax caudimaculatus</i> (Lucena, 1987)	0	2	2	0.04	4.5	35827
<i>Gnathocharax steindachneri</i> Fowler, 1913	0	8	8	0.15	4.5	35887
<i>Hemigrammus belottii</i> (Steindachner, 1882)	0	349	349	6.34	27.3	35888
<i>Hemigrammus gracilis</i> (Lütken, 1875)	1	0	1	0.02	4.5	35806
<i>Hemigrammus ocellifer</i> (Steindachner, 1882)	179	42	221	4.01	31.8	35788, 35847, 35889
<i>Hemigrammus schmardae</i> (Steindachner, 1882)	0	380	380	6.90	18.2	35829, 35875
<i>Hemigrammus vorderwinkleri</i> Géry, 1963	649	0	649	11.78	27.3	35766, 35792
<i>Heterocharax</i> sp. (juvenile)	0	1	1	0.02	4.5	35900
<i>Hyphessobrycon cf. tukunai</i> Géry, 1965	0	16	16	0.29	9.1	35849, 35876
<i>Hyphessobrycon heterorhabdus</i> (Ulrey, 1894)	763	362	1125	20.42	72.7	35769, 35807, 35864
<i>Iguanodectes geisleri</i> Géry, 1970	7	3	10	0.18	9.1	35808
<i>Iguanodectes variatus</i> Géry, 1993	0	9	9	0.16	18.2	35832, 35865, 35891
<i>Moenkhausia mikia</i> Marinho and Langeani, 2010	0	1	1	0.02	4.5	35893
<i>Moenkhausia chrysargyrea</i> (Günther, 1864)	1	0	1	0.02	4.5	35815
<i>Moenkhausia cotinho</i> Eigenmann, 1908	0	2	2	0.04	9.1	35894, 35910
<i>Moenkhausia cf. diktyota</i> Lima and Toledo-Piza, 2001	0	12	12	0.22	9.1	35855, 35863
<i>Moenkhausia oligolepis</i> (Günther, 1864)	7	0	7	0.13	4.5	35791

TABLE 1. CONTINUED.

TAXON	BASINS			FR (%)	FO (%)	INPA #
	PURUS	MADEIRA	N			
<i>Priocharax ariel</i> Weitzman and Vari, 1987	0	173	173	3.14	18.2	35825
<i>Tyttocharax madeirae</i> Fowler, 1913	12	33	45	0.82	13.6	35811
Crenuchidae						
<i>Crenuchus spilurus</i> Günther, 1863	0	93	93	1.69	36.4	35871
<i>Elachocharax junki</i> (Géry, 1971)	0	29	29	0.53	13.6	35885
<i>Elachocharax pulcher</i> Myers, 1927	0	1	1	0.02	4.5	35845
<i>Microcharacidium weitzmani</i> Buckup, 1993	6	126	132	2.40	36.4	35774, 35833, 35877
<i>Odontocharacidium aphanes</i> (Weitzman and Kanazawa, 1977)	0	31	31	0.56	22.7	35823, 35834, 35896
Curimatidae						
<i>Cyphocharax pantostictos</i> Vari and Barriga S., 1990	1	0	1	0.02	4.5	35813
Erythrinidae						
<i>Erythrinus erythrinus</i> (Bloch and Schneider, 1801)	13	18	31	0.56	63.6	35767, 35821, 35854, 35914, 35920
<i>Hoplias malabaricus</i> (Bloch, 1794)	1	8	9	0.16	31.8	35814, 35830, 35848, 35890, 35922
<i>Hoplerythrinus unitaeniatus</i> (Spix and Agassiz, 1829)	7	0	7	0.13	22.7	35768
Gasteropelecidae						
<i>Carnegiella marthae</i> Myers, 1927	0	1	1	0.02	4.5	35883
<i>Carnegiella strigata</i> (Günther, 1864)	19	38	57	1.03	45.5	35906
Lebiasinidae						
<i>Copella nattereri</i> (Steindachner, 1876)	0	623	623	11.31	50.0	35820, 35838, 35844, 35860, 35907, 35919
<i>Copella nigrofasciata</i> (Meinken, 1952)	80	0	80	1.45	22.7	35778
<i>Nannostomus eques</i> Steindachner, 1876	0	1	1	0.02	4.5	35895
<i>Pyrrhulina brevis</i> Steindachner, 1876	63	62	125	2.27	72.7	35818, 35824, 35835, 35841, 35851, 35857, 35867, 35879, 35911, 35924
<i>Pyrrhulina zigzag</i> Zarske and Géry, 1997	170	0	170	3.09	27.3	35794
CYPRINODONTIFORMES						
Rivulidae						
<i>Rivulus kirovskyi</i> Costa, 2004	22	3	25	0.45	27.3	35775, 35852, 35869
<i>Rivulus</i> sp.	85	36	121	2.20	68.2	35770
GYMNOTIFORMES						
Gymnotidae						
<i>Gymnotus carapo</i> Linnaeus, 1758	3	3	6	0.11	13.6	35787, 35804, 35872
<i>Gymnotus coatesi</i> La Monte, 1935	0	1	1	0.02	4.5	35921
<i>Gymnotus coropinae</i> Hoedeman, 1962	19	41	60	1.09	63.6	35797, 35861, 35873, 35916
<i>Gymnotus</i> sp.1	2	1	3	0.05	13.6	35771, 35805, 35917
<i>Gymnotus</i> sp.2	0	15	15	0.27	9.1	35846, 35862
Hypopomidae						
<i>Brachyhypopomus brevirostris</i> (Steindachner, 1868)	0	5	5	0.09	9.1	35843, 35904
<i>Brachyhypopomus</i> sp.1	0	54	54	0.98	18.2	35882, 35905, 35913
<i>Brachyhypopomus</i> sp.2	19	1	20	0.36	31.8	35777, 35785, 35793, 35796, 35870
<i>Brachyhypopomus</i> sp.3	1	0	1	0.02	4.5	35816
<i>Hypopygus lepturus</i> Hoedeman, 1962	14	89	103	1.87	40.9	35789, 35831, 35840, 35923
<i>Microsternarchus bilineatus</i> Fernández-Yépez, 1968	0	28	28	0.51	18.2	35850
<i>Microsternarchus</i> sp.	2	46	48	0.87	22.7	35856, 35866, 35878
<i>Steatogenys duidae</i> (La Monte, 1929)	0	1	1	0.02	4.5	35826
Rhamphichthyidae						
<i>Gymnorhamphichthys petiti</i> Géry and Vu-Tân-Tuê, 1964	1	0	1	0.02	4.5	35803
Sternopygidae						
<i>Sternopygus aequilabiatus</i> (Humboldt, 1805)	0	1	1	0.02	4.5	35836
PERCIFORMES						
Cichlidae						
<i>Acaronia nassa</i> (Heckel, 1840)	0	1	1	0.02	4.5	35902
<i>Aequidens</i> sp.1	6	2	8	0.15	22.7	35781, 35842
<i>Aequidens</i> sp.2	0	2	2	0.04	4.5	35859
<i>Apistogramma agassizi</i> (Steindachner, 1875)	67	149	216	3.92	77.3	35782, 35881, 35903

TABLE 1. CONTINUED.

TAXON	BASINS			FR (%)	FO (%)	INPA #
	PURUS	MADEIRA	N			
<i>Apistogramma regani</i> Kullander, 1980	15	0	15	0.27	9.1	35783
<i>Crenicara punctulatum</i> (Günther, 1863)	0	1	1	0.02	4.5	35884
<i>Crenicichla inpa</i> Ploeg, 1991	1	0	1	0.02	4.5	35812
Gobiidae						
<i>Microphilypinus ternetzi</i> Myers, 1927	0	18	18	0.33	9.1	35892
SILURIFORMES						
Auchenipteridae						
<i>Tatia brunnea</i> Mees, 1974	0	1	1	0.02	4.5	35880
<i>Tatia gyrina</i> (Eigenmann and Allen, 1942)	1	3	4	0.07	13.6	35799, 35899, 35912
Callichthyidae						
<i>Megalechis picta</i> (Müller and Troschel, 1849)	3	6	9	0.16	18.2	35790, 35909
Cetopsidae						
<i>Helogenes marmoratus</i> Günther, 1863	10	40	50	0.91	50.0	35779, 35822, 35828, 35839, 35874
Doradidae						
<i>Physopyxis ananas</i> Sousa and Rapp Py-Daniel, 2005	0	11	11	0.20	4.5	35897
Heptapteridae						
<i>Gladioganis conquistador</i> Lundberg, Bornbusch and Mago-Leccia, 1991	26	109	135	2.45	45.5	35786, 35802, 35886, 35908, 35915
<i>Rhamdia quelen</i> (Quoy and Gaimard, 1824)	0	3	3	0.05	13.6	35858, 35868, 35898
Loricariidae						
<i>Ancistrus</i> sp.	2	0	2	0.04	4.5	35800
<i>Otocinclus caxarari</i> Schaefer, 1997	4	0	4	0.07	4.5	35809
Pseudopimelodidae						
<i>Batrochoglanis raninus</i> (Valenciennes, 1840)	1	0	1	0.02	4.5	35773
Trichomycteridae						
<i>Ituglanis amazonicus</i> (Steindachner, 1882)	6	0	6	0.11	13.6	35780
<i>Trichomycterus hasemani</i> (Eigenmann, 1914)	0	5	5	0.09	9.1	35901
SYNBRANCHIFORMES						
Synbranchidae						
<i>Synbranchus madeirae</i> Rosen and Rumney, 1972	2	0	2	0.04	9.1	35795, 35817
<i>Synbranchus</i> sp.	2	20	22	0.40	40.9	35798, 35810, 35853, 35918, 35837
N = 78	2332	3176	5508			

ACKNOWLEDGMENTS: We thank the GEOMA Project for financial and logistical support of the field and laboratory work at INPA. We thank Mario Cohn-Haft for coordinating the project and the Petrobras for providing a helicopter that greatly eased the access to the study area. The first author acknowledges the Instituto Internacional de Educação do Brasil (IEB) – BECA Program for the Masters scholarship, and the Environmental Sciences Postgraduate Program of the UFPA-MPEG-EMBRAPA. JZ receives a productivity grant from CNPq (process # 307464/2009-1). IBAMA provided the license to collect biological material. Two anonymous reviewers made valuable contributions to improve the manuscript.

LITERATURE CITED

- American Veterinary Association Medical Association. 2001. Report of the AVMA panel on euthanasia. *JAVMA* 218: 669-696.
- Araújo-Lima, C.A.R.M., L.F. Jiménez, R.S. Oliveira, P.C. Eterovick, U. Mendonza and A. Jerolimki. 1999. Relação entre o número de espécies de peixes, complexidade de hábitat e ordem do riacho nas cabeceiras de um tributário do rio Urubu, Amazônia Central. *Acta Limnológica Brasiliensis* 11(2): 127-135.
- Barthem, R.B. 2004. Aquatic biota; p. 62-79 In J.P.R. Capobianco, A. Veríssimo, A. Moreira, D. Sawyer, I. Santos and L.P. Pinto (ed.). *Biodiversity in the Brazilian Amazon: assessment and priority actions for conservation, sustainable use and benefit sharing*. São Paulo: Estação Liberdade and Instituto Socioambiental.
- Böhlke, J.E., S.H. Weitzman and N.A. Menezes. 1978. Estado atual da sistemática dos peixes de água doce da América do Sul. *Acta Amazonica* 8(4): 657-677.
- Buckup, P.A. 1993. Review of the characidiin fishes (Teleostei: Characiformes), with descriptions of four new genera and ten new species. *Ichthyological Exploration of Freshwaters* 4(2): 97-154.
- Buckup, P.A., N.A. Menezes and M.S. Ghazzi. 2007. *Catálogo das espécies de peixes de água doce do Brasil*. Rio de Janeiro: Museu Nacional, Universidade Federal do Rio de Janeiro, Série Livros 23. 195 p.
- Bührnheim, C.M. and C. Cox-Fernandes. 2001. Low seasonal variation of fish assemblages in Amazonian rain forest streams. *Ichthyological Exploration of Freshwaters* 12(1): 65-78.
- Burgess, W.E. 1989. *An atlas of freshwater and marine catfishes*. Neptune: TFH Publications. 784 p.
- Castro, R.M.C. 1999. Evolução da ictiofauna de riachos sul-americanos: padrões gerais e possíveis processos causais; p. 139-155 In E.P. Caramaschi, R. Mazzoni and P.R. Peres-Neto (ed.). *Ecologia de peixes de riachos*. Rio de Janeiro: PPGE-UFRRJ (Série Oecologia Brasiliensis vol. VI).
- Costa, W.J.E.M. 2004. *Rivulus kirovskyi*, a new killifish from the central Amazon, Brazil (Cyprinodontiformes: Rivulidae). *Neotropical Ichthyology* 2(1): 9-12.
- Crampton, W.G.R. and J.S. Albert. 2003. Redescription of *Gymnotus coropinae* (Gymnotiformes, Gymnotidae), an often misidentified species of Neotropical electric fish, with notes on natural history and electric signals. *Zootaxa* 348: 1-20.
- Crampton, W.G.R. and J.S. Albert. 2004. Redescription of *Gymnotus coatesi* (Gymnotiformes, Gymnotidae), a Rare Species of Electric Fish from the Lowland Amazon Basin, with Descriptions of Osteology, Electric Signals and Ecology. *Copeia* 2004(3): 525-533.
- Crampton, W.G.R., D.H. Thorsen and J.S. Albert. 2005. Three new species from a diverse, sympatric assemblage of the electric fish *Gymnotus* (Gymnotiformes: Gymnotidae) in the lowland Amazon Basin, with notes on ecology. *Copeia* 2005(1): 82-99.

- Crampton, W.G.R., K.G. Hulen and J.S. Albert. 2004. *Sternopygus branco*: A new species of neotropical electric fish (Gymnotiformes: Sternopygidae) from the lowland Amazon basin, with descriptions of osteology, ecology, and electric organ discharges. *Copeia* 2004(2): 245-259.
- Crampton, W.G.R., N.R. Lovejoy and J.S. Albert. 2003. *Gymnotus ucumari*: a new species of Neotropical electric fish from the Peruvian Amazon (Ostariophysi: Gymnotidae), with notes on ecology and electric organ discharges. *Zootaxa* 277: 1-18.
- Dias, M.S., W.E. Magnusson and J. Zuanon. 2009. Effects of Reduced-Impact Logging on Fish Assemblages in Central Amazonia. *Conservation Biology* 24(1): 278-286.
- Espírito-Santo, H.M.V., W.E. Magnusson, J. Zuanon, F.P. Mendonça and V.L. Landeiro. 2009. Seasonal variation in the composition of fish assemblages in small Amazonian forest streams: evidence for predictable changes. *Freshwater Biology* 54(3): 536-548.
- Favorito, S.E., A.M. Zanata and M.I. Assumpção. 2005. A new *Synbranchus* (Teleostei: Synbranchiformes: Synbranchidae) from Ilha de Marajó, Pará, Brazil, with notes on its reproductive biology and larval development. *Neotropical Ichthyology* 3(3): 319-328.
- Fearnside, P.M. and P.M.L.A. Graça. 2006. BR-319: Brazil's Manaus-Porto Velho Highway and the potential impact of linking the arc of deforestation to central Amazonia. *Environmental Management* 38(5): 705-716.
- Fink, W.I. and S.V. Fink. 1978. A Amazônia Central e seus peixes. *Acta Amazonica* (Suplemento) 8(4): 19-42.
- Géry, J. 1977. *Characoids of the World*. Neptune: TFH Publications. 772 p.
- Géry, J. 1993. Description de trois espèces nouvelles du genre *Iguanodectes* (Pisces, Characiformes, Characidae), avec quelques données récentes sur les autres espèces. *Revue française d'Aquariologie* 19(4): 97-105.
- Huber, J.H. 1992. *Rivulus: Ecobiogeography - Relationships*. *Laboratoire d'Ichtyologie general et appliquée*. Paris: Museum National d'Histoire Naturelle. 572 p.
- Kullander, S.O. 1986. *Cichlid fishes of the Amazon River drainage of Peru*. Stockholm: Swedish Museum of Natural History. 431 p.
- Kullander, S.O. 1989. Description of a new *Acaronia* species from the Rio Orinoco and Rio Negro drainages. *Zoologica Scripta* 18(3): 447-452.
- Kullander, S.O. 1995. Three new cichlid species from southern Amazonia: *Aequidens gerciliae*, *A. epae* and *A. michaeli*. *Ichthyological Exploration of Freshwaters* 6:149-170.
- Kullander, S.O. and E.J.G. Ferreira. 1991. A new *Aequidens* species from the Rio Trombetas, Brasil, and redescription of *Aequidens pallidus*. *Zoologica Scripta* 19(4): 425-433.
- Lima, F.C.T. and M. Toledo-Piza. 2001. New species of *Moenkhausia* (Characiformes: Characidae) from the Rio Negro of Brazil. *Copeia* 2001(4): 1058-1063.
- Lowe-McConnell, R.H. 1999. *Estudos ecológicos de comunidades de peixes tropicais*. São Paulo: Editora da Universidade de São Paulo. 534 p.
- Lucena, C.A.S. 1987. Revisão e redefinição do gênero Neotropical *Charax* Scopoli, 1777, com a descrição de quatro espécies novas (Pisces; Characiformes; Characidae). *Comunicações do Museu de Ciências da PUCRS* 40: 5-124.
- Lundberg, J.G. 2005. *Gymnorhamphichthys bogardusi*, a new species of sand knifefish (Gymnotiformes: Rhamphichthyidae) from the Rio Orinoco, South America. *Annals of the South African Museum* 479: 1-4.
- Mago-Leccia, F. 1994. *Electric fishes of the continental waters of America*. Caracas: Fundacion para el Desarrollo de las Ciencias Fisicas, Matematicas y Naturales. 206 p.
- Marinho, M.M.F. and F. Langeani. 2010. A new species of *Moenkhausia* from the rio Amazonas and rio Orinoco basins (Characiformes: Characidae). *Zootaxa* 2577:57-68.
- Mendonça, F.P., W.E. Magnusson and J. Zuanon. 2005. Relationships between habitat characteristics and fish assemblages in small streams of central Amazonia. *Copeia* 2005(4): 751-764.
- Nogueira, C., P.A. Buckup, N.A. Menezes, O.T. Oyakawa, T.P. Kasecker, M.B. Ramos Neto and J.M.C. da Silva. 2010. Restricted-range fishes and the conservation of Brazilian freshwaters. *PLoS One* 5(6): e11390.
- Oliveira R.R., M.S. Rocha, M.B. Anjos, J. Zuanon and L.H. Rapp Py-Daniel. 2009. Fish fauna of small streams of the Catua-Ipixuna Extractive Reserve, State of Amazonas, Brazil. *Check List* 5(2): 154-172.
- Oyakawa, O.T. and G.M.T. Mattox. 2009. Revision of the Neotropical trahiras of the *Hoplias lacerdae* species-group (Ostariophysi: Characiformes: Erythrinidae) with descriptions of two new species. *Neotropical Ichthyology* 7(2): 117-140.
- Peres, C.A. 2005. Why we need megareserves in Amazonia. *Conservation Biology* 19(3): 728-733.
- Peres, C.A., T.A. Gardner, J. Barlow, J. Zuanon, F. Michalski, A.C. Lees, I.C.G. Vieira, F.M.S. Moreira and K.J. Feeley. 2010. Biodiversity conservation in human-modified Amazonian forest landscapes. *Biological Conservation* 143 (2010) 2314-2327.
- Radambrasil. 1978. *Mineral, D. N. P. Folha SB. 20 Purus: geologia, geomorfologia, pedologia, vegetação e uso potencial da terra*. Rio de Janeiro: Instituto Brasileiro de Geografia e Estatística, 566 p.
- Rapp Py-Daniel, L.H. and E.L.M. Leão. 1991. A coleção de peixes do INPA: base do conhecimento científico sobre a ictiofauna amazônica gerado pelo Instituto Nacional de Pesquisas da Amazônia; p. 299-312. In A.L. Val, R. Figliuolo and E. Feldberg (ed.). *Bases científicas para estratégias de preservação e desenvolvimento da Amazônia: fatos e perspectivas*. Manaus: Instituto Nacional de Pesquisas da Amazônia.
- Reis, R.E. 1997. Revision of the Neotropical genus *Hoplosternum* (Ostariophysi: Siluriformes: Callichthyidae) with the description of two new genera and three new species. *Ichthyological Exploration of Freshwaters* 7: 299-326.
- Reis, R.E., P-Y. Le Bail and J.H.A. Mol. 2005. New arrangement in the synonymy of *Megalechis* Reis, 1997 (Siluriformes: Callichthyidae). *Copeia* 2005(3): 678-682.
- Reis, R.E., S.O. Kullander and C.J. Ferraris Jr. 2003. *Check list of the freshwater fishes of South and Central America*. Porto Alegre: EDIPUCRS. 742 p.
- Rocha, M.S., R.R. de Oliveira and L.H.R. Py-Daniel. 2008. A new species of *Gladioglanis* Ferraris and Mago-Leccia from rio Aripuanã, Amazonas, Brazil (Siluriformes: Heptapteridae). *Neotropical Ichthyology* 6(3): 433-438.
- Römer, U. 2002. *Cichlid Atlas, Volume 1. Natural History of South American Dwarf Cichlids. Parts 1 and 2*. Germany: Mergus, Melle. 1311 p.
- Sabino, J. and J. Zuanon. 1998. A stream fish assemblage in Central Amazônia: distribution, activity pattern and feeding behaviour. *Ichthyological Exploration of Freshwaters* 8(3): 201-210.
- Sarmento-Soares, L.M. and R.F. Martins-Pinheiro. 2008. A systematic revision of *Tatia* (Siluriformes: Auchenipteridae: Centromochlinae). *Neotropical Ichthyology* 6(3): 495-542.
- Schaefer, S.A. 1997. The Neotropical cascudinhos: Systematics and biogeography of the *Otocinclus* catfishes (Siluriformes: Loricariidae). *Proceedings of the Academy of Natural Sciences of Philadelphia* 148: 1-120.
- Silfvergrip, A.M.C. 1996. *A systematic revision of the Neotropical catfish genus Rhamdia (Teleostei, Pimelodidae)*. Stockholm: Swedish Museum of Natural History. 156 p.
- Silva, C.P.D. 1995. Community structure of fish in urban and natural streams in the Central Amazon. *Amazoniana* 13(3/4): 221-236.
- Sousa, L.M. and L.H.R. Py-Daniel. 2005. Description of two new species of *Physopyxis* and redescription of *P. lyra* (Siluriformes: Doradidae). *Neotropical Ichthyology* 3(4): 625-636.
- Thacker, C.E., F. Pezold and R.D. Suttkus. 2006. Redescription of the dwarf neotropical eleotrid genus *Leptophilypnus* (Teleostei: Gobioidi), including a new species and comments on *Microphilypnus*. *Copeia* 2006(3): 489-499.
- Vari, R.P. 1992. Systematics of the Neotropical Characiform genus *Cyphocharax* Fowler (Pisces, Ostariophysi). *Smithsonian Contributions to Zoology* 529: 1-137.
- Vari, R.P. and H. Ortega. 1986. The catfishes of the Neotropical family Helogenidae (Ostariophysi: Siluroidei). *Smithsonian Contributions to Zoology* 442: 1-20.
- Weitzman, S.H. 1960. Further notes on the relationships and classification of the South American characid fishes of the subfamily Gasteropelecinae. *Stanford Ichthyological Bulletin* 7(4): 217-239.
- Weitzman, S.H. 1978. Three new species of fishes of the genus *Nannostomus* from the Brazilian states of Pará and Amazonas (Teleostei: Lebiasinidae). *Smithsonian Contributions to Zoology* 263: 1-14.
- Weitzman, S.H. and R.P. Vari. 1987. Two new species and a new genus of miniature characid fishes (Teleostei: Characiformes) from northern South America. *Proceedings of The Biological Society of Washington* 100(3): 640-652.
- Zarske, A. and J. Géry. 1997. Ein neuer Salmmleraus Peru. *Das Aquarium* 336: 12-17.
- Zarske, A. and J. Géry. 2006. Zur identität von *Copella nattereri* (Steindachner, 1876) einschließlich der Beschreibung einer neuen Art (Teleostei: Characiformes: Lebiasinidae). *Zoologische Abhandlungen (Dresden)* 56:15-46.

RECEIVED: February 2011

LAST REVISED: October 2011

ACCEPTED: October 2011

PUBLISHED ONLINE: December 2011

EDITORIAL RESPONSIBILITY: Rubens Pazza